arXiv:1407.5433v3 [math.NT] 10 Jul 2015

ARITHMETIC PROPERTIES OF PARTITION TRIPLES WITH ODD PARTS DISTINCT

LIUQUAN WANG

Abstract. Let $\mathrm{pod}_{-3}(n)$ denote the number of partition triples of n where the odd parts in each partition are distinct. We find many arithmetic properties of $\mathrm{pod}_{-3}(n)$ involving the following infinite family of congruences: for any integers $\alpha \geq 1$ and $n \geq 0$,

$$\mathrm{pod}_{-3}\Big(3^{2\alpha+2}n + \frac{23 \times 3^{2\alpha+1}+3}{8}\Big) \equiv 0 \pmod 9.$$

We also establish some arithmetic relations between $\mathrm{pod}(n)$ and $\mathrm{pod}_{-3}(n)$, as well as some congruences for $\mathrm{pod}_{-3}(n)$ modulo 7 and 11.

1. Introduction

Let $\mathrm{pod}_{-k}(n)$ be defined by the generating function

$$\sum_{n=0}^{\infty} \mathrm{pod}_{-k}(n)q^n = \frac{(-q;q^2)_\infty^k}{(q^2;q^2)_\infty^k} = \frac{1}{\psi(-q)^k},$$

where $\psi(q)$ is one of Ramanujan's theta functions, namely

$$\psi(q) = \sum_{n=0}^{\infty} q^{n(n+1)/2} = \frac{(q^2;q^2)_\infty^2}{(q;q)_\infty}.$$

The combinatorial interpretation is that $\mathrm{pod}_{-k}(n)$ denotes the number of partition k-tuples of n where in each partition the odd parts are distinct. For $k = 1$, $\mathrm{pod}_{-1}(n)$ is often denoted as $\mathrm{pod}(n)$.

In 2010, Hirschhorn and Sellers [5] investigated the arithmetic properties of $\mathrm{pod}(n)$. They obtained some interesting congruences. For example, for integers $\alpha \geq 0$ and $n \geq 0$, they showed that

$$\mathrm{pod}\Big(3^{2\alpha+3}n + \frac{23 \times 3^{2\alpha+2}+1}{8}\Big) \equiv 0 \pmod 3.$$

Some internal congruences were also found, such as

$$\mathrm{pod}(81n+17) \equiv 5\,\mathrm{pod}(9n+2) \pmod{27}.$$

In 2011, using the tool of modular forms, Radu and Sellers [8] obtained other deep congruences for $\mathrm{pod}(n)$ modulo 5 and 7, such as

$$\mathrm{pod}(135n+8) \equiv \mathrm{pod}(135n+107) \equiv \mathrm{pod}(135n+116) \equiv 0 \pmod 5.$$

$$\mathrm{pod}(567n+260) \equiv \mathrm{pod}(567n+449) \equiv 0 \pmod 7.$$

Chen and Xia [3] followed by studying the arithmetic properties of $\mathrm{pod}_{-2}(n)$. They found two infinite families of congruences modulo 3 and 5, respectively. For more properties of $\mathrm{pod}(n)$, see [7, 10].

2010 *Mathematics Subject Classification.* Primary 05A17; Secondary 11P83.

Key words and phrases. Congruences; partition triples; distinct odd parts; theta functions.

This paper appears in International Journal of Number Theory, doi:10.1142/S1793042115500773.

Inspired by their work, we study the arithmetic properties of $\mathrm{pod}_{-3}(n)$, the number of partition triples of n where the odd parts in each partition are distinct. In Section 2, we will present an infinite family of Ramanujan-type congruences: for any integers $\alpha \geq 1$ and $n \geq 0$,

$$\mathrm{pod}_{-3}\Big(3^{2\alpha+2}n + \frac{23\times 3^{2\alpha+1}+3}{8}\Big) \equiv 0 \pmod 9.$$

In Section 3, we establish the following arithmetic relations between $\mathrm{pod}(n)$ and $\mathrm{pod}_{-3}(n)$:

$$\mathrm{pod}_{-3}(3n+2) \equiv 6\,\mathrm{pod}(9n+5) \pmod 9,$$
$$\mathrm{pod}_{-3}(3n+1) \equiv 3\,\mathrm{pod}(9n+2) \pmod{18}.$$

In Section 4, we will prove

$$\mathrm{pod}_{-3}(7n+3) \equiv (-1)^{n+1} r_3(8n+3) \pmod 7,$$
$$\mathrm{pod}_{-3}(11n+10) \equiv 4(-1)^n r_7(8n+7) \pmod{11},$$

where $r_k(n)$ denotes the number of representations of n as sum of k squares. Many congruences will also be deduced from these two relations.

2. An Infinite Family of Congruences Modulo 9

Before our discussion, we observe that (see [3, 5])

$$\psi(q) = 1+q+q^3+q^6+q^{10}+q^{15}+\cdots = A(q^3)+q\psi(q^9).$$

where

$$A(q) = \frac{(q^2;q^2)_\infty (q^3;q^3)_\infty^2}{(q;q)_\infty (q^6;q^6)_\infty}.$$

The following lemma plays a key role in both [3] and [5], which will also be used frequently in our paper.

Lemma 2.1. *(Cf.* [5, Lemma 2.1]*.) We have*

$$A(q^3)^3 + q^3\psi(q^9)^3 = \frac{\psi(q^3)^4}{\psi(q^9)}$$

and

$$\frac{1}{\psi(q)} = \frac{\psi(q^9)}{\psi(q^3)^4}\left(A(q^3)^2 - qA(q^3)\psi(q^9) + q^2\psi(q^9)^2\right).$$

For convenience, throughout the paper, we denote $s = A(q^3)$ and $t = \psi(q^9)$. Then Lemma 2.1 can be rewritten as

$$s^3 + q^3t^3 = \frac{\psi(q^3)^4}{\psi(q^9)}, \tag{2.1}$$

$$\frac{1}{\psi(q)} = \frac{\psi(q^9)}{\psi(q^3)^4}(s^2 - qst + q^2t^2). \tag{2.2}$$

To establish some congruences using q series, we also need the following fact.

Lemma 2.2. *(Cf.* [8, Lemma 1.2]*.) Let p be a prime and α be a positive integer. Then*

$$(q;q)_\infty^{p^\alpha} \equiv (q^p;q^p)_\infty^{p^{\alpha-1}} \pmod{p^\alpha}.$$

Since $\psi(q) = \frac{(q^2;q^2)_\infty^2}{(q;q)_\infty}$, we deduce from Lemma 2.2 that $\psi(q)^p \equiv \psi(q^p) \pmod p$ for any prime p. We will use this fact many times.

We are going to establish an infinite family of congruences modulo 9.

Theorem 2.1. *We have*

$$\sum_{n=0}^{\infty}(-1)^n \mathrm{pod}_{-3}(3n)q^n \equiv \frac{\psi(q^3)}{\psi(q)^4} \pmod 9.$$

$$\sum_{n=0}^{\infty}(-1)^n \mathrm{pod}_{-3}(3n+1)q^n \equiv 3\psi(q)A(q)^2 \pmod 9.$$

$$\sum_{n=0}^{\infty}(-1)^n \mathrm{pod}_{-3}(3n+2)q^n \equiv -3\psi(q^3)\psi(q)A(q) \pmod 9.$$

Proof. By (2.2) we have

$$\sum_{n=0}^{\infty}(-1)^n \mathrm{pod}_{-3}(n)q^n = \frac{1}{\psi(q)^3} = \frac{\psi(q^9)^3}{\psi(q^3)^{12}}(s^2-qst+q^2t^2)^3. \tag{2.3}$$

Note that

$$(s^2-qst+q^2t^2)^3 = (s^6-7q^3s^3t^3+q^6t^6)+q(-3s^5t+6q^3s^2t^4)+q^2(6s^4t^2-3q^3st^5). \tag{2.4}$$

Applying (2.1) we obtain

$$\begin{aligned}\sum_{n=0}^{\infty}(-1)^{3n} \mathrm{pod}_{-3}(3n)q^{3n} &= \frac{\psi(q^9)^3}{\psi(q^3)^{12}}(s^6-7q^3s^3t^3+q^6t^6)\\ &\equiv \frac{\psi(q^9)^3}{\psi(q^3)^{12}}(s^3+q^3t^3)^2\\ &\equiv \frac{\psi(q^9)}{\psi(q^3)^4} \pmod 9.\end{aligned}$$

Replacing q^3 by q, we get the first congruence identity.

Similarly,

$$\begin{aligned}\sum_{n=0}^{\infty}(-1)^{3n+1} \mathrm{pod}_{-3}(3n+1)q^{3n+1} &= q\frac{\psi(q^9)^3}{\psi(q^3)^{12}}\big(-3s^2t(s^3+q^3t^3)+9q^3s^2t^4\big)\\ &\equiv -3q\frac{\psi(q^9)^3}{\psi(q^3)^{12}}s^2t(s^3+q^3t^3)\\ &\equiv -3q\frac{\psi(q^9)^3}{\psi(q^3)^8}A(q^3)^2 \pmod 9.\end{aligned}$$

Dividing both sides by $-q$, then replacing q^3 by q and applying Lemma 2.2, we obtain

$$\sum_{n=0}^{\infty}(-1)^n \mathrm{pod}_{-3}(3n+1)q^n \equiv 3\frac{\psi(q^3)^3}{\psi(q)^8}A(q)^2 \equiv 3\psi(q)A(q)^2 \pmod 9.$$

This completes the proof of the second congruence identity.

In the same way, we have

$$\begin{aligned}\sum_{n=0}^{\infty}(-1)^{3n+2} \mathrm{pod}_{-3}(3n+2)q^{3n+2} &= q^2\frac{\psi(q^9)^3}{\psi(q^3)^{12}}\big(9s^4t^2-3st^2(s^3+q^3t^3)\big)\\ &\equiv -3q^2\frac{\psi(q^9)^3}{\psi(q^3)^{12}}st^2(s^3+q^3t^3)\\ &\equiv -3q^2\frac{\psi(q^9)^4}{\psi(q^3)^8}A(q^3) \pmod 9.\end{aligned}$$

Dividing both sides by q^2, then replacing q^3 by q and applying Lemma 2.2, we obtain

$$\sum_{n=0}^{\infty}(-1)^n \mathrm{pod}_{-3}(3n+2)q^n \equiv -3\frac{\psi(q^3)^4}{\psi(q)^8}A(q) \equiv -3\psi(q^3)\psi(q)A(q) \pmod 9.$$

□

Theorem 2.2. *For any integers $\alpha \geq 1$ and $n \geq 0$, we have*

$$\sum_{n=0}^{\infty}(-1)^n \mathrm{pod}_{-3}\Big(3^{2\alpha}n + \frac{5\times 3^{2\alpha}+3}{8}\Big)q^n \equiv (-1)^{\alpha-1}\frac{\psi(q^3)^4}{\psi(q)^7} \pmod 9 \tag{2.5}$$

and

$$\sum_{n=0}^{\infty}(-1)^n \mathrm{pod}_{-3}\Big(3^{2\alpha+1}n + \frac{7\times 3^{2\alpha+1}+3}{8}\Big)q^n \equiv (-1)^{\alpha-1}\frac{\psi(q^3)^5}{\psi(q)^8} \pmod 9. \tag{2.6}$$

Proof. By (2.2) and the first congruence in Theorem 2.1, we have

$$\sum_{n=0}^{\infty}(-1)^n \mathrm{pod}_{-3}(3n)q^n \equiv \frac{\psi(q^3)}{\psi(q)^4} = \frac{\psi(q^9)^4}{\psi(q^3)^{15}}(s^2-qst+q^2t^2)^4 \pmod 9. \tag{2.7}$$

Note that

$$\begin{aligned}(s^2-qst+q^2t^2)^4 = {}& (s^8-16q^3s^5t^3+10q^6s^2t^6)+q(-4s^7t+19q^3s^4t^4-4q^6st^7)\\ &+q^2(10s^6t^2-16q^3s^3t^5+q^6t^8).\end{aligned} \tag{2.8}$$

Collecting all the terms of the form q^{3k+2} in (2.7), we obtain

$$\begin{aligned}\sum_{n=0}^{\infty}(-1)^{3n+2}\mathrm{pod}_{-3}(9n+6)q^{3n+2} &= q^2\frac{\psi(q^9)^4}{\psi(q^3)^{15}}(10s^6t^2-16q^3s^3t^5+q^6t^8)\\ &\equiv q^2\frac{\psi(q^9)^4}{\psi(q^3)^{15}}t^2(s^3+q^3t^3)^2\\ &\equiv q^2\frac{\psi(q^9)^4}{\psi(q^3)^7} \pmod 9.\end{aligned}$$

Dividing both sides by q^2 and then replacing q^3 by q, we get exactly (2.5) with $\alpha=1$.

Now suppose we already have (2.5) for a given α. By (2.2) it follows that

$$\frac{\psi(q^3)^4}{\psi(q)^7} = \frac{\psi(q^9)^7}{\psi(q^3)^{24}}(s^2-qst+q^2t^2)^7.$$

Collecting all the terms of the form q^{3k+2} in $(s^2-qst+q^2t^2)^7$, we obtain

$$\begin{aligned}&q^2(28s^{12}t^2-266q^3s^9t^5+357q^6s^6t^8-77q^9s^3t^{11}+q^{12}t^{14})\\ &\equiv q^2t^2(s^{12}+4q^3s^9t^3+6s^6q^6t^6+4q^9s^3t^9+q^{12}t^{12})\\ &\equiv q^2t^2(s^3+q^3t^3)^4 \pmod 9.\end{aligned}$$

Hence when collecting all the terms of the form q^{3k+2} in (2.5) and applying (2.1), we get

$$\sum_{n=0}^{\infty}(-1)^{3n+2}\mathrm{pod}_{-3}\Big(3^{2\alpha}(3n+2)+\frac{5\times 3^{2\alpha}+3}{8}\Big)q^{3n+2} \equiv (-1)^{\alpha-1}q^2\frac{\psi(q^9)^5}{\psi(q^3)^8} \pmod 9.$$

Dividing both sides by q^2 and then replacing q^3 by q, we deduce that

$$\sum_{n=0}^{\infty}(-1)^n \mathrm{pod}_{-3}\Big(3^{2\alpha+1}n + \frac{7\times 3^{2\alpha+1}+3}{8}\Big)q^n \equiv (-1)^{\alpha-1}\frac{\psi(q^3)^5}{\psi(q)^8} \pmod 9. \tag{2.9}$$

Again by (2.2) we have

$$\frac{\psi(q^3)^5}{\psi(q)^8}=\frac{\psi(q^9)^8}{\psi(q^3)^{27}}(s^2-qst+q^2t^2)^8.$$

Collecting all the terms of the form q^{3k+1} in $(s^2-qst+q^2t^2)^8$, we get

$$\begin{aligned}&q(-8s^{15}t+266q^3s^{12}t^4-1016q^6s^9t^7+784q^9s^6t^{10}-112q^{12}s^3t^{13}+q^{15}t^{16})\\&\equiv qt(s^{15}+5q^3s^{12}t^3+10q^6s^9t^6+10q^9s^6t^9+5q^{12}s^3t^{12}+q^{15}t^{15})\\&\equiv qt(s^3+q^3t^3)^5 \pmod 9.\end{aligned}$$

If we collect all the terms of the form q^{3k+1} in (2.9), we obtain

$$\sum_{n=0}^{\infty}(-1)^{3n+1}\mathrm{pod}_{-3}\Big(3^{2\alpha+1}(3n+1)+\frac{7\times 3^{2\alpha+1}+3}{8}\Big)q^{3n+1}\equiv(-1)^{\alpha-1}q\frac{\psi(q^9)^4}{\psi(q^3)^7} \pmod 9.$$

Now dividing both sides by $-q$ and then replacing q^3 by q, we get

$$\sum_{n=0}^{\infty}(-1)^{n}\mathrm{pod}_{-3}\Big(3^{2\alpha+2}n+\frac{5\times 3^{2\alpha+2}+3}{8}\Big)q^{n}=(-1)^{(\alpha+1)-1}\frac{\psi(q^3)^4}{\psi(q)^7} \pmod 9.$$

By doing induction on α, we complete our proof. □

Corollary 2.1. *For any integer $n\geq 0$, we have*

$$\begin{aligned}\mathrm{pod}_{-3}(81n+51)&\equiv-\mathrm{pod}_{-3}(9n+6) \pmod 9,\\ \mathrm{pod}_{-3}(27n+24)&\equiv-\mathrm{pod}_{-3}(243n+213) \pmod 9.\end{aligned}$$

Proof. Let $\alpha=1,2$ in (2.5). We obtain

$$\sum_{n=0}^{\infty}(-1)^n\mathrm{pod}_{-3}(9n+6)q^n\equiv\frac{\psi(q^3)^4}{\psi(q)^7}\equiv-\sum_{n=0}^{\infty}(-1)^n\mathrm{pod}_{-3}(81n+51)q^n \pmod 9.$$

The first congruence follows immediately.

The second congruence can be obtained in a similar way by letting $\alpha=1,2$ in (2.6). □

Lemma 2.3. *The coefficients of q^{3n+2} in the series expansion of $\frac{1}{\psi(q)^8}$ is divisible by 9.*

Proof. By (2.2) we have

$$\frac{1}{\psi(q)^8}=\frac{\psi(q^9)^8}{\psi(q^3)^{32}}\big(s^2-qst+q^2t^2\big)^8.$$

Collecting all the terms of the form q^{3k+2} in $(s^2-qst+q^2t^2)^8$, we obtain

$$9q^2(4s^{14}t^2-56q^3s^{11}t^5+123q^6s^8t^8-56q^9s^5t^{11}+4q^{12}s^2t^{14}).$$

It is obvious that all the coefficients are divisible by 9. □

Now we are able to prove the following theorem.

Theorem 2.3. *For any integers $\alpha\geq 1$ and $n\geq 0$, we have*

$$\mathrm{pod}_{-3}\Big(3^{2\alpha+2}n+\frac{23\times 3^{2\alpha+1}+3}{8}\Big)\equiv 0 \pmod 9.$$

Proof. By Lemma 2.3, we know the coefficient of the term q^{3n+2} in the left hand side of (2.6) is divisible by 9. Hence we have

$$\mathrm{pod}_{-3}\Big(3^{2\alpha+1}(3n+2)+\frac{7\times 3^{2\alpha+1}+3}{8}\Big)\equiv 0 \pmod 9.$$

This is the desired result we want. □

Note that for different integers $\alpha \geq 1$, the arithmetic sequences $\left\{3^{2\alpha+2}n + \frac{23\times 3^{2\alpha+1}+3}{8} : n = 0, 1, 2, \cdots\right\}$ mentioned in Theorem 2.3 are disjoint, and they account for

$$\frac{1}{3^4} + \frac{1}{3^6} + \cdots = \frac{1}{72}$$

of all nonnegative integers. Thus we have the following corollary.

Corollary 2.2. $\mathrm{pod}_{-3}(n)$ *is divisible by* 9 *for at least* 1/72 *of all nonnegative integers.*

3. Arithmetic Relations Between $\mathrm{pod}(n)$ and $\mathrm{pod}_{-3}(n)$

There are some nontrivial arithmetic relations between $\mathrm{pod}(n)$ and $\mathrm{pod}_{-3}(n)$. First we recall some congruence identities discovered in [5].

Lemma 3.1. *(Cf.* [5, Section 3]*.)* $\sum\limits_{n=0}^{\infty}(-1)^n \mathrm{pod}(3n+2)q^n = \frac{\psi(q^3)^3}{\psi(q)^4}$.

Lemma 3.2. *(Cf.* [5, Section 4]*.)*

$$\sum_{n=0}^{\infty}(-1)^n \mathrm{pod}(9n+2)q^n \equiv A(q)^2\frac{\psi(q^3)^2}{\psi(q)^5} \pmod 9,$$
$$\sum_{n=0}^{\infty}(-1)^n \mathrm{pod}(9n+5)q^n \equiv 4A(q)\frac{\psi(q^3)^3}{\psi(q)^5} \pmod 9,$$
$$\sum_{n=0}^{\infty}(-1)^n \mathrm{pod}(9n+8)q^n \equiv \frac{\psi(q^3)^4}{\psi(q)^5} \pmod 9.$$

Theorem 3.1. *For any integer $n \geq 0$, we have*

$$\mathrm{pod}_{-3}(3n+2) \equiv 6\,\mathrm{pod}(9n+5) \pmod 9,$$
$$\mathrm{pod}_{-3}(3n+1) \equiv 3\,\mathrm{pod}(9n+2) \pmod{18}.$$

Proof. By Lemma 2.2, we have $\psi(q^3)^2 \equiv \psi(q)^6 \pmod 3$. Hence

$$24A(q)\frac{\psi(q^3)^3}{\psi(q)^5} \equiv -3\psi(q^3)\psi(q)A(q) \pmod 9.$$

Hence by the second identity in Lemma 3.2 and the third identity in Theorem 2.1, we get

$$\mathrm{pod}_{-3}(3n+2) \equiv 6\,\mathrm{pod}(9n+5) \pmod 9.$$

Now we turn to the proof of the second congruence. Since $\psi(q^3)^2 \equiv \psi(q)^6 \pmod 3$, we have

$$3A(q)^2\frac{\psi(q^3)^2}{\psi(q)^5} \equiv 3A(q)^2\psi(q) \pmod 9.$$

Hence by the first identity in Lemma 3.2 and the second identity in Theorem 2.1, we obtain

$$\mathrm{pod}_{-3}(3n+1) \equiv 3\,\mathrm{pod}(9n+2) \pmod 9.$$

The remaining task is to show that $\mathrm{pod}_{-3}(3n+1) \equiv \mathrm{pod}(9n+2) \pmod 2$.

By Lemma 3.1 and (2.2), we have

$$\sum_{n=0}^{\infty}(-1)^n \mathrm{pod}(3n+2)q^n = \frac{\psi(q^3)^3}{\psi(q)^4} = \frac{\psi(q^9)^4}{\psi(q^3)^{13}}(s^2 - qst + q^2t^2)^4. \tag{3.1}$$

Collecting all the terms of the form q^{3n} in the right hand side of (3.1), by (2.8) we obtain

$$\frac{\psi(q^9)^4}{\psi(q^3)^{13}}(s^8 - 16q^3s^5t^3 + 10q^6s^2t^6) \equiv \frac{\psi(q^9)^4}{\psi(q^3)^{13}}A(q^3)^8 \pmod 2.$$

Hence we have

$$\sum_{n=0}^{\infty}(-1)^{3n}\mathrm{pod}(9n+2)q^{3n}\equiv\frac{\psi(q^9)^4}{\psi(q^3)^{13}}A(q^3)^8\pmod 2.$$

Replacing q^3 by q, we get

$$\sum_{n=0}^{\infty}(-1)^{n}\mathrm{pod}(9n+2)q^{n}\equiv\frac{\psi(q^3)^4}{\psi(q)^{13}}A(q)^8\pmod 2. \tag{3.2}$$

Back to the expression of $\sum_{n=0}^{\infty}(-1)^n\mathrm{pod}_{-3}(n)q^n$ in (2.3), from (2.4) we have

$$\begin{aligned}\sum_{n=0}^{\infty}(-1)^{3n+1}\mathrm{pod}_{-3}(3n+1)q^{3n+1}&=q\frac{\psi(q^9)^3}{\psi(q^3)^{12}}(-3s^5t+6q^3s^2t^4)\\&\equiv q\frac{\psi(q^9)^4}{\psi(q^3)^{12}}A(q^3)^5\pmod 2.\end{aligned}$$

Now dividing both sides by $-q$ and replacing q^3 by q, we get

$$\sum_{n=0}^{\infty}(-1)^{n}\mathrm{pod}_{-3}(3n+1)q^{n}\equiv\frac{\psi(q^3)^4}{\psi(q)^{12}}A(q)^5\pmod 2. \tag{3.3}$$

By Lemma 2.2, we have $(q^2;q^2)_\infty\equiv(q;q)_\infty^2\pmod 2$ and $(q^3;q^3)_\infty^2\equiv(q^6;q^6)_\infty\pmod 2$. Hence

$$A(q)=\frac{(q^2;q^2)_\infty(q^3;q^3)_\infty^2}{(q;q)_\infty(q^6;q^6)_\infty}\equiv(q;q)_\infty\pmod 2,$$

and

$$\psi(q)=\frac{(q^2;q^2)_\infty^2}{(q;q)_\infty}\equiv(q;q)_\infty^3\pmod 2.$$

Therefore, we have $A(q)^3\equiv\psi(q)\pmod 2$. Thus

$$\frac{\psi(q^3)^4}{\psi(q)^{13}}A(q)^8\equiv\frac{\psi(q^3)^4}{\psi(q)^{12}}A(q)^5\pmod 2.$$

From (3.2), (3.3) and the congruence above, we deduce that $\mathrm{pod}_{-3}(3n+1)\equiv\mathrm{pod}(9n+2)\pmod 2$, which completes our proof. □

4. Congruences Modulo 7 And 11

For nonnegative integers n and k, let $r_k(n)$ denote the number of representations of n as sum of k squares, and $t_k(n)$ denote the number of representations of n as sum of k triangular numbers. Let $\sigma(n)$ denote the sum of positive divisors of n.

The method used in this section is quite similar to the one used in [9], where the author applied the properties of $r_k(n)$ to give new congruences satisfied by overpartition function.

Lemma 4.1. *For any prime $p\geq 3$, we have*

$$t_4\Big(pn+\frac{p-1}{2}\Big)\equiv t_4(n)\pmod p,\quad t_8(pn+p-1)\equiv t_8(n)\pmod p.$$

Proof. From Theorem 3.6.3 in [2], we know $t_4(n)=\sigma(2n+1)$. For any positive integer N, we have

$$\sigma(N)=\sum_{d|N,\,p|d}d+\sum_{d|N,\,p\nmid d}d\equiv\sum_{d|N,\,p\nmid d}d\pmod p.$$

Let $N=2n+1$ and $N=p(2n+1)$ respectively. It is easy to see that $\sigma(p(2n+1))\equiv\sigma(2n+1)\pmod p$, which clearly implies the first congruence.

From equation (3.8.3) in page 81 of [2], we know

$$t_8(n)=\sum_{\substack{d|(n+1)\\ d\,\mathrm{odd}}}\Big(\frac{n+1}{d}\Big)^3.$$

Hence similarly we can prove the second congruence. □

Lemma 4.2. *(Cf. [1].) For $1\leq k\leq 7$, we have*

$$r_k(8n+k)=2^k\Big(1+\frac{1}{2}\binom{k}{4}\Big)t_k(n).$$

Theorem 4.1. *For any integer $n\geq 0$, we have*

$$\mathrm{pod}_{-3}(7n+3)\equiv(-1)^{n+1}r_3(8n+3)\pmod 7.$$

Proof. Let $p=7$ in Lemma 4.1. We deduce that $t_4(7n+3)\equiv t_4(n)\pmod 7$.

From definition we have

$$\psi(q)^7\sum_{n=0}^{\infty}\mathrm{pod}_{-3}(n)(-q)^n=\psi(q)^4=\sum_{n=0}^{\infty}t_4(n)q^n.$$

By Lemma 2.2, we have $\psi(q)^7\equiv\psi(q^7)\pmod 7$. Replacing $\psi(q)^7$ by $\psi(q^7)$ modulo 7 and extracting those terms of the form q^{7n+3} in the above identity, we obtain

$$\psi(q^7)\sum_{n=0}^{\infty}\mathrm{pod}_{-3}(7n+3)(-q)^{7n+3}\equiv\sum_{n=0}^{\infty}t_4(7n+3)q^{7n+3}\pmod 7.$$

Dividing both sides by $-q^3$, then replacing q^7 by q and replacing $t_4(7n+3)$ by $t_4(n)$, we get

$$\sum_{n=0}^{\infty}\mathrm{pod}_{-3}(7n+3)(-q)^n\equiv-\psi(q)^3\equiv-\sum_{n=0}^{\infty}t_3(n)q^n\pmod 7.$$

Hence we have $\mathrm{pod}_{-3}(7n+3)\equiv(-1)^{n+1}t_3(n)\pmod 7$.

Let $k=3$ in Lemma 4.2. We obtain $t_3(n)=r_3(8n+3)/8$, from which the theorem follows. □

We can deduce some interesting congruences from this theorem.

Lemma 4.3. *(Cf. [6].) Let $p\geq 3$ be a prime. For any integers $n\geq 1$ and $\alpha\geq 0$, we have*

$$r_3(p^{2\alpha}n)=\left(\frac{p^{\alpha+1}-1}{p-1}-\Big(\frac{-n}{p}\Big)\frac{p^\alpha-1}{p-1}\right)r_3(n)-p\frac{p^\alpha-1}{p-1}r_3(n/p^2).$$

Here $(\frac{\cdot}{p})$ denotes the Legendre symbol, and we take $r_3(n/p^2)=0$ unless $p^2|n$.

Theorem 4.2. *Let p be a prime and $p\equiv 6\pmod 7$. Let N be a positive integer which is coprime to p and $p^3N\equiv 3\pmod 8$. We have*

$$\mathrm{pod}_{-3}\Big(\frac{7p^3N+3}{8}\Big)\equiv 0\pmod 7.$$

Proof. Let $\alpha=1$ and $n=pN$ in Lemma 4.3. We have

$$r_3(p^3N)=(p+1)r_3(pN)\equiv 0\pmod 7.$$

Let $n=\frac{p^3N-3}{8}$. By Theorem 4.1, we deduce that

$$\mathrm{pod}_{-3}\Big(\frac{7p^3N+3}{8}\Big)=\mathrm{pod}_{-3}(7n+3)\equiv(-1)^{n+1}r_3(p^3N)\equiv 0\pmod 7.$$

□

For example, let $p=13$ and $N=8n+7$, where $n\not\equiv 4 \pmod{13}$ is a nonnegative integer. We establish a Ramanujan-type congruence as follows:

$$\mathrm{pod}_{-3}(15379n+13457)\equiv 0 \pmod 7.$$

Theorem 4.3. *Let $p\geq 3$ be a prime, N a positive integer which is coprime to p and $pN\equiv 3 \pmod 8$. Let α be any nonnegative integer.*
(1) If $p\equiv 1 \pmod 7$, then

$$\mathrm{pod}_{-3}\Big(\frac{7p^{14\alpha+13}N+3}{8}\Big)\equiv 0 \pmod 7.$$

(2) If $p\not\equiv 0$ or $1 \pmod 7$, then

$$\mathrm{pod}_{-3}\Big(\frac{7p^{12\alpha+11}N+3}{8}\Big)\equiv 0 \pmod 7.$$

Proof. (1) Let $n=pN$ in Lemma 4.3, and then we replace α by $7\alpha+6$. Since $p\equiv 1 \pmod 7$, we have

$$\frac{p^{7\alpha+7}-1}{p-1}=1+p+\cdots+p^{7\alpha+6}\equiv 0 \pmod 7.$$

Hence $r_3(p^{14\alpha+13}N)\equiv 0 \pmod 7$.

Let $n=\frac{p^{14\alpha+13}N-3}{8}$. By Theorem 4.1, we deduce that

$$\mathrm{pod}_{-3}\Big(\frac{7p^{14\alpha+13}N+3}{8}\Big)=\mathrm{pod}_{-3}(7n+3)\equiv(-1)^{n+1}r_3(p^{14\alpha+13}N)\equiv 0 \pmod 7.$$

(2) Let $n=pN$ in Lemma 4.3, and then we replace α by $6\alpha+5$. Note that $p^{6\alpha+6}\equiv 1 \pmod 7$, we get $r_3(p^{12\alpha+11}N)\equiv 0 \pmod 7$.

Let $n=\frac{p^{12\alpha+11}N-3}{8}$. From Theorem 4.1 we obtain

$$\mathrm{pod}_{-3}\Big(\frac{7p^{12\alpha+11}N+3}{8}\Big)=\mathrm{pod}_{-3}(7n+3)\equiv(-1)^{n+1}r_3(p^{12\alpha+11}N)\equiv 0 \pmod 7.$$

□

Theorem 4.4. *Let $a\in\{3,19,27\}$. For any integers $n\geq 0$ and $\alpha\geq 1$, we have*

$$\mathrm{pod}_{-3}\Big(7^{2\alpha+2}n+\frac{a\cdot 7^{2\alpha+1}+3}{8}\Big)\equiv 0 \pmod 7.$$

Proof. Let $p=7$ and $n=7m+r$ ($r\in\{3,5,6\}$) in Lemma 4.3. It is easy to deduce that $r_3(7^{2\alpha}(7m+r))\equiv 0 \pmod 7$.

Let $n=7^{2\alpha+1}m+\frac{a\cdot 7^{2\alpha}-3}{8}$ ($a\in\{3,19,27\}$) in Theorem 4.1. We have

$$\mathrm{pod}_{-3}\Big(7^{2\alpha+2}m+\frac{a\cdot 7^{2\alpha+1}+3}{8}\Big)=\mathrm{pod}_{-3}(7n+3)\equiv(-1)^{n+1}r_3(8n+3) \pmod 7.$$

Since $r_3(8n+3)=r_3(7^{2\alpha}(56m+a))\equiv 0 \pmod 7$, we complete our proof. □

Note that for different pairs (α,a), $\alpha\geq 1$ and $a\in\{3,19,27\}$, the arithmetic sequences $\big\{7^{2\alpha+2}n+\frac{a\cdot 7^{2\alpha+1}+3}{8}: n=0,1,2,\cdots\big\}$ are disjoint, and they account for

$$3\Big(\frac{1}{7^4}+\frac{1}{7^6}+\cdots+\frac{1}{7^{2\alpha+2}}+\cdots\Big)=\frac{1}{784}$$

of all nonnegative integers. Thus we have

Corollary 4.1. $\mathrm{pod}_{-3}(n)$ *is divisible by* 7 *for at least* $1/784$ *of all nonnegative integers.*

Theorem 4.5. *For any integers $n\geq 0$, we have*

$$\mathrm{pod}_{-3}(11n+10)\equiv 4(-1)^n r_7(8n+7) \pmod{11}.$$

Proof. Let $p=11$ in Lemma 4.1. We deduce that $t_8(11n+10)\equiv t_8(n) \pmod{11}$. By Lemma 2.2 we have $\psi(q)^{11}\equiv\psi(q^{11}) \pmod{11}$. Hence

$$\psi(q^{11})\sum_{n=0}^{\infty}\mathrm{pod}_{-3}(n)(-q)^n\equiv\psi(q)^8=\sum_{n=0}^{\infty}t_8(n)q^n \pmod{11}.$$

If we extract all the terms of the form q^{11n+10}, we obtain

$$\psi(q^{11})\sum_{n=0}^{\infty}\mathrm{pod}_{-3}(11n+10)(-q)^{11n+10}\equiv\sum_{n=0}^{\infty}t_8(11n+10)q^{11n+10} \pmod{11}.$$

Dividing both sides by q^{10} and replacing q^{11} by q, we have

$$\psi(q)\sum_{n=0}^{\infty}\mathrm{pod}_{-3}(11n+10)(-q)^n\equiv\sum_{n=0}^{\infty}t_8(11n+10)q^n\equiv\sum_{n=0}^{\infty}t_8(n)q^n=\psi(q)^8 \pmod{11}.$$

Therefore,

$$\sum_{n=0}^{\infty}\mathrm{pod}_{-3}(11n+10)(-q)^n\equiv\psi(q)^7=\sum_{n=0}^{\infty}t_7(n)q^n \pmod{11}.$$

Comparing the coefficients of q^n on both sides, we obtain

$$\mathrm{pod}_{-3}(11n+10)\equiv(-1)^n t_7(n) \pmod{11}.$$

Let $k=7$ in Lemma 4.2. We deduce that $t_7(n)=\frac{1}{2368}r_7(8n+7)\equiv 4r_7(8n+7) \pmod{11}$. The theorem follows. □

Lemma 4.4. *(Cf. [4].) Let $p\geq 3$ be a prime, and n a nonnegative integer such that $p^2\nmid n$. For any integer $\alpha\geq 0$, we have*

$$r_7(p^{2\alpha}n)=\left(\frac{p^{5\alpha+5}-1}{p^5-1}-p^2\big(\frac{-n}{p}\big)\frac{p^{5\alpha}-1}{p^5-1}\right)r_7(n).$$

Theorem 4.6. *Let $p\geq 3$ be a prime and $\alpha\geq 0$ be an integer. Suppose $1\leq r\leq 7$ and $pr\equiv 7 \pmod 8$, $0\leq a<p$ and $p\nmid 8a+r$, where both a and r are integers.*
(1) If $p\equiv 1,3,4,5,9 \pmod{11}$, then

$$\mathrm{pod}_{-3}\Big(11\cdot p^{22\alpha+22}n+\frac{(88a+11r)p^{22\alpha+21}+3}{8}\Big)\equiv 0 \pmod{11}.$$

(2) If $p\equiv 2,6,7,8,10 \pmod{11}$, then

$$\mathrm{pod}_{-3}\Big(11\cdot p^{4\alpha+4}n+\frac{(88a+11r)p^{4\alpha+3}+3}{8}\Big)\equiv 0 \pmod{11}.$$

Proof. (1) Replacing α by $11\alpha+10$ and setting $n=pN$ in Lemma 4.4, where N is coprime to p. Note that $p\equiv 1,3,4,5,9 \pmod{11}$ implies $p^5\equiv 1 \pmod{11}$, we have

$$\frac{p^{55\alpha+55}-1}{p^5-1}=1+p^5+p^{10}+\cdots+p^{5(11\alpha+10)}\equiv 0 \pmod{11}.$$

From which we know $r_7(p^{22\alpha+21}N)\equiv 0 \pmod{11}$.

Let $n=p^{22\alpha+22}m+\frac{(8a+r)p^{22\alpha+21}-7}{8}$. We have

$$r_7(8n+7)=r_7\big(p^{22\alpha+21}(8pm+8a+r)\big)\equiv 0 \pmod{11}.$$

By Theorem 4.5, we obtain

$$\mathrm{pod}_{-3}\Big(11\cdot p^{22\alpha+22}m+\frac{(88a+11r)p^{22\alpha+21}+3}{8}\Big)=\mathrm{pod}_{-3}(11n+10)\equiv 0 \pmod{11}.$$

(2) Let $n = pN$, where N is coprime to p, and we replace α by $2\alpha+1$ in Lemma 4.4. Note that $p \equiv 2, 6, 7, 8, 10 \pmod{11}$ implies $p^5 \equiv -1 \pmod{11}$, we have $p^{10\alpha+10} \equiv 1 \pmod{11}$. We deduce that $r_7(p^{4\alpha+3}N) \equiv 0 \pmod{11}$.

Let $n = p^{4\alpha+4}m + \frac{(8a+r)p^{4\alpha+3}-7}{8}$. We have

$$r_7(8n+7) = r_7\big(p^{4\alpha+3}(8pm+8a+r)\big) \equiv 0 \pmod{11}.$$

By Theorem 4.5, we obtain

$$\mathrm{pod}_{-3}\Big(11 \cdot p^{4\alpha+4}m + \frac{(88a+11r)p^{4\alpha+3}+3}{8}\Big) = \mathrm{pod}_{-3}(11n+10) \equiv 0 \pmod{11}.$$

□

For example, let $p = 7$ in part (2) of Theorem 4.6. Accordingly we set $r = 1$ and $a \in \{0, 1, 2, 3, 4, 5\}$. For any integer $n \geq 0$, we have

$$\mathrm{pod}_{-3}\Big(11 \cdot 7^{4\alpha+4}n + \frac{(88a+11) \cdot 7^{4\alpha+3}+3}{8}\Big) \equiv 0 \pmod{11}.$$

Starting from the above congruence, we can prove the following corollary by a way similar to the proof of Corollary 4.1,

Corollary 4.2. $\mathrm{pod}_{-3}(n)$ *is divisible by* 11 *for at least* $1/4400$ *of all nonnegative integers.*

References

[1] P. Barrucand, S. Cooper, and M. D. Hirschhorn, Relations between squares and triangles, Discrete Math. **248** (2002), 245–247.

[2] B. C. Berndt, *Number Theory in the Spirit of Ramanujan*, Am. Math. Soc. (Providence, 2006).

[3] W. Y. C. Chen and B. L. S. Lin, Congruences for bipartitions with odd parts distinct, Ramanujan J. **25** (2) (2011), 277–293.

[4] S. Cooper, Sums of five, seven and nine squares, Ramanujan J. **6** (2002), 469–490.

[5] M. D. Hirschhorn and J. A. Sellers, Arithmetic properties of partitions with odd parts distinct, Ramanujan J. **22** (2010), 273–284.

[6] M. D. Hirschhorn and J. A. Sellers, On representations of a number as a sum of three squares, Discrete Math. **199** (1999), 85–101.

[7] J. Lovejoy and R. Osburn, Quadratic forms and four partition functions modulo 3, Integers **11** (2011), A4.

[8] S. Radu and J. A. Sellers, Congruence properties modulo 5 and 7 for the pod function, Int. J. Number Theory **07** (2011), 2249–2259.

[9] L. Wang, Another proof of a conjecture of Hirschhorn and Sellers on overpartitions, J. Integer Seq. **17** (2014), Article 14.9.8, 8pp.

[10] L. Wang, New congruances of partitions where the old parts are distinct, J. Integer Seq. 18 (2015) Article 15.4.2, 8 pp.

Department of Mathematics, National University of Singapore, Singapore, 119076, SINGAPORE

Department of Mathematics, Zhejiang University, Hangzhou, 310027, CHINA

E-mail address: mathlqwang@163.com; wangliuquan@u.nus.edu